# 2

## **POCUMENTATION PAGE**

Form Approved OMB No 0704-0188

				·			
	U CLMSS	HELATION		16 RESTRICTIVE		•	
2a SECURITY CLASSIFICATION AUTHORITY NA			3 DISTRIBUTION AVAILABILITY OF REPORT				
26 DECLASSIFICATION DOWNGRADING SCHEDULE NA			Distribution Unlimited				
4 PERFORMING ORGANIZATION REPORT NUMBER(S)			5 MONITORING ORGANIZATION REPORT NUMBERIS				
				NA	1		
6a NAME OF	PERFORMING	ORGANIZATION	6b OFFICE SYMBOL (If applicable)	78 NAME OF MO	ONITORING DRG	AN 741-0N	6. C. 19. 30
The University of Connecticut NA			NA	Office of Naval Research			
Molecula 354 Mans Storrs,	sfield Roa CT 06269	Biology, U-13 ad 9-2131			incy St. , VA 2221	7-5000	
8a NAME OF ORGANIZA	FUNDING 'SPO	ONSORING	8b OFFICE SYMBOL (If applicable)	9 PROCUREMENT	CINSTRUMENT	DENT FICATION N	∪ <b>M</b> BEF
	of Naval [		ONR	N00014-88			
	City, State, and			10 SOURCE OF F		ــــــــــــــــــــــــــــــــــــــ	
	800 N. Quincy St. Arlington, VA 22217-5000			PROGRAM ELEMENT NO 61153N	PROJECT NO RRO4106	1412-034	WORK UNIT ACCESSION NO
11 TITLE (Inci	ude Security C	lassification)		<b>4</b> ,	<u> </u>	<del></del>	<del></del>
		nd Fate of Bac	cterial Sulfona	tes			
12 PERSONA. Leadbett		rd R., Godchau	ıx, W., III				
13a TYPE OF		136 TIME CO	DVERED	14 DATE OF REPO		Day) 15 PACE	<i>C</i> 0.≯*
Annual	NTARY NOTAT		10 10 _12/00	1989, Janu	ary US	13	
17	'*A2O)	<del></del>	18 SUBJECT TERMS (	Continue on reverse	e if necessary ar	nd identify by blo	(k number)
FIELD	GROUP	SUB-GROUP	sulfonates; b				n;
00	03	<del></del>	biocorrosion; biotransforma	sulfonate r tion:(κ→) →	eduction;	bacterial;	
19 ABSTRACT	(Continue on	reverse if necessary	and identify by block n				
resear sulfor and no isethican be the su unknow	ch project nates by b on-phototr onate, ar utilized onate-s vn. Plans	t focused on bacteria. Describes on the control ones cand taurine. It is a second to the second of t	gress made during an examination scribed are find an metabolize so an some instance of one or another find year effort	of the form dings that boot old be sulfo es the entire bacterium ate of the r	ation and oth photot nates such e organosu, in other emainder o ject are o	degradation rophic bact as cysteat lfur compou instances f the molecutlined.	of eria e, nd only
WUNCLAS	SIFIED UNLIMIT	ED SAME AS R	PT DTIC USERS	(U)	nclude Area Cod	(e) 2.16 OF 16E 5	rMao.
M. Marr			Previous editions are	202-696-4		ONR CLASSICATION	CH THIS FACE (U)
	J, 2014 00		. ICTOUTEUROUNDE			= = : : : : :	

S/N 0102-LF-014-6603





ANNUAL REPORT:

CONTRACT N00014-88-K-0120

R&T CODE 4412-034

PRINCIPAL INVESTIGATORS:

War ACT

Edward R. Leadbetter, Walter Godchauv III

CONTRACTOR:

The University of Connecticut, Storrs

CONTRACT TITLE:

Formation and Fate of Bacterial Sulfonates

START DATE:

1 January 1988

DATE OF REPORT:

5 JANUARY 1989

RESEARCH OBJECTIVES: a. to isolate and characterize bacteria able (i) to utilize sulfonate-sulfur for biosynthesis or (ii) to carry out dissimilative sulfonate reduction, then to assess the physiology and biochemistry of these two different processes. b. to examine the routes of sulfonate-S formation in the simple gliding bacteria that are known to biosynthesize the sulfonate, cysteate, in significant quantities.

PROGRESS:

a. Examination of the biotransformation (biodegradation) of sulfonates has been focused on two physiological types of bacteria—those phototrophic bacteria able to live "facultatively" either by phototrophy in anaerobic environments or by light-independent respiration under aerobic conditions, as well as non-phototrophic (chemoorganotrophic) bacteria that are strictly respiratory; in this latter category anaerobic respiration with nitrate has also been examined. Thus far the bulk of the bacteria isolated have been from fresh-water and soil habitats.

PHOTOTROPHS. Rhodobacters and rhodopseudomonads (Rhodospirillaceae) have been obtained by enrichment culturing [in the light, anaerobically] using sulfonates (e.g., cysteate, isethionate, taurine) as sole sources of S, and the ability of the bacteria to utilize sulfonate-S for assimilative purposes has been compared under the two growth conditions—phototrophic and respiratory. Several salient facts emerge: sulfonate-S and sulfate-S support equal amounts of growth [thiosulfate did not serve effectively]; a nutritional specificity, or selectiveness, exists in some strains (i.e., not all three of the sulfonates can serve as S sources), while other strains are nutritionally versatile; under phototrophic conditions many, but not all, strains are able to utilize the entire sulfonate molecule as sole source of C, S, and N. This is the first demonstration of the biotransformation of sulfonates under phototrophic, anaerobic conditions!

Three cyanobacteria—a strain each of <u>Synechococcus</u>, <u>Anabena</u>, and <u>Nostoc</u>—have been examined for their ability to utilize the sulfonate taurine as sole source of S for their oxygenic phototrophic growth; only <u>Anabena</u> and <u>Nostoc</u> were able to do so, and the latter could do so when grown either phototrophically or heterotrophically.

CHEMORGANOTROPHS. Pseudomonads and other aerobic bacteria. The enrichment culture and isolation of chemoorganotrophs able to utilize sulfonates as sole source of C, S, E (energy) and (where applicable) N has led to the demonstration that the isolates are able to attack and utilize taurine as sole C, N, S, and energy source, and that the majority of the strains also are able to so utilize isethionate [excepting N utilization]; strangely, only one isolate is able to degrade cysteate.

### b. Cysteate synthesis and its regulation.

Although on comparative biochemical grounds one would expect the biosynthesis of cysteate to proceed from the amino acid cysteine, via an oxidation of the sulfhydryl group to a sulfinate, thence to the sulfonate moiety, our earlier studies demonstrating that the carbon of cysteine did not appear in cysteate indicated that this expected route was not followed in Cytophaga johnsonae. Isolation of a mutant unable to synthesize cysteine, but still able to form sulfonolipids of which cysteate is the "head-group" precursor, has led to a quite clear demonstration that the biosynthesis of cysteate proceeds by a pathway other than one involving cyteine's carbons. A number of possible precursors are being evaluated experimentally.

As one means of establishing the pathway for cysteate synthesis, a kinetic analysis of sulfate uptake and its incorporation into low molecular weight intracellular metabolites was undertaken. Using wild-type C. johnsonae, we have demonstrated that cysteate is indeed a major S-containing metabolite in these bacteria and, in addition that these organisms possess regulatory features for sulfate uptake and assimilation that seem quite unlike those studied in enteric, and other, bacteria and in plants and thus may possess a novel pattern of regulation of sulfur metabolism.

WORK PLAN (YEAR 2).

a. PHOTOTROPHS and AEROBIC CHEMOORGANOTROPHS. The identification of the bacteria already isolated should be completed, as should also the isolation and characterization of marine isolates. Thus for both bacterial groups then, it may be possible to begin focusing on the nature of the enzyme(s) involved in cleaving the C-S bond of sulfonates, and determining whether the enzymes involved are identical in those instances where the S is used only for assimilative purposes as contrasted with the cases where the entire sulfonate molecule is degraded.

SULFONATE RESPIRATION. Although stable enrichment cultures that seemingly effect the reduction of sulfonates (the only known electron acceptor added) are in hand, we have had no luck yet in obtaining pure cultures that carry out this transformation(s). Efforts to do so will continue, using more stringent anaerobic techniques than have heretofore been employed by us.

b. CYSTEATE BIOSYNTHESIS. During year two we expect to have obtained definitive evidence supporting, or ruling out, the involvement of several prospective precursor metabolites of cysteate; aspartate is the molecule that, on the basis of present experimental evidence, seems the most likely candidate. Once evidence from study of intact cells is indicative of the likely biosynthetic pathway we will turn to examination of cell-extracts in order to provide firm evidence of the route for biosynthesis.

#### PUBLICATIONS, REPORTS.

- 1. Eysteine is not an obligatory intermediate in the biosynthesis of cysteate by Cytophaga johnsonae. E. F. Gilmore. W. Godohaux III., E. R. Leadbetter, Biochem. Biophys. Res. Commun., SUBMITTED
- 2. Regulation of dysteate synthesis in the gliding bacterium <u>Sytophaga nohnsonae</u>. D. F. Gilmore, W. Goddhaux, E. R. Leadbetter. SuPMITTED for presentation at the Annual Meeting, Amer. Soc. Microbiol., 1989.
- 3. Bacterial utilization of sulforates. S. M. Green, A. F. Seitz, E. F. Leadbetter. SUBMITTED for presentation, Annual Mesting, Amer. Soc. Microbiol., 1992
- 4. A progress report covering fix months' research was prepared and mulled, in June 1988, to those on the distribution list.

TRAINING ACTIVITIES. Graduate students Catherine Greene, David Gilmore. Thomas Fitta, Lisa Gorski and Maria Uria have been supported in whole on part by this contract. An undergraduate student, Angelica Seitz, has worked on this project, and will pentimus to be so during at least the first six months of year two, or a part time basis.

Designathic data requested: Women, three graduate, 1 undergraduate student: Non-citizens: 1 (M. unia, hative of Bolivia).

#### AWARDE/FELLOWSHIPE:

ERL was an Inter-Adagen. Exchange Fellow of the E. S. and Mungarian National Adagenies of Science for three months in Fall 1929; site was at the Biological Peseanon Senter. Spaged, Hungary where dyamobacterial stress responses and sulforate utilization were studied.

Acces	sion For	
NTIS	GRA&I	Y
DTIC	TAB	A
Unann	ounced	$\Box$
Justi	fication	
D		
Ву		
Distr	ibution/	
Avai	lobility	Codes
	Av.il and	/or
Dist	Special	
	1	
A 1		
Zi-li		



# Distribution List for Annual and Final Reports

- 1. Put a cover page (Form DD 1473) on your report and attach a copy of the distribution list. Mail one copy of the report to each person on the contractor subset list attached on which your name appears. The other subset list is for your information only. Please don't forget to attach this distribution list to your report otherwise the folks below think they have mistakenly received the copy meant for the Molecular Biology Program and forward it to us.
- Mail two copies to (include a DTIC Form 50 with these two copies too)
   Administrator
   Defense Technical Information Center
   Building 5, Cameron Station
   Alexandria, VA 22314
- 3. Mail one copy to each of the following:
- (a) Dr. Michael Marron
  ONR Code 1141
  Molecular Biology Program
  800 N. Quincy Street
  Arlington, VA 22217-5000
- (b) Administrative Contracting Officer
  ONR Resident Representative
  (address varies see copy of your grant)
- (c) Director,
  Applied Research Directorate
  ONR Code 12
  800 N. Quincy Street
  Arlington, VA 22217-5000
- (d) Director
  Office of Naval Technology
  Code 22
  800 N. Quincy Street
  Arlington, VA 22217-5000

- (e) Director
  Chemical and Biological Sci Div
  Army Research Office
  P. 0. Box 12211
  Research Triangle Park, NC 27709
- (f) Life Sciences Directorate
  Air Force Office of Scientific Research
  Bolling Air Force Base
  Washington, DC 20332
- (g) Director
  Naval Research Laboratory
  Technical Information Div, Code 2627
  Washington, DC 20375